


CLAIMS

WHAT IS CLAIMED IS:

5 1. A method of treating affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other
 10 feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and
 15 spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in
 20 mammals comprising administering to the mammal a therapeutically effective amount of a compound of Formula (1):



(1)

25

and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and

pharmaceutically acceptable salt forms thereof,
wherein:

Ar is selected from phenyl, naphthyl, pyridyl,
5 pyrimidinyl, triazinyl, furanyl, thieryl,
benzothienyl, benzofuranyl, 2,3-
dihydrobenzofuranyl, 2,3-dihydrobenzothienyl,
indanyl, 1,2-benzopyranyl, 3,4-dihydro-1,2-
benzopyranyl, tetrinalinyl, each Ar optionally
10 substituted with 1 to 5 R⁴ groups and each Ar is
attached to an unsaturated carbon atom;

R¹ is independently selected at each occurrence from
H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl,
15 halo, CN, C₁-C₄ haloalkyl, C₁-C₁₂ hydroxyalkyl,
C₂-C₁₂ alkoxyalkyl, C₂-C₁₀ cyanoalkyl, C₃-C₆
cycloalkyl, C₄-C₁₀ cycloalkylalkyl, NR⁹R¹⁰, C₁-
C₄ alkyl-NR⁹R¹⁰, NR⁹COR¹⁰, OR¹¹, SH or S(O)_nR¹²;

20 R² is selected from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-
C₄ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₀
cycloalkylalkyl, C₁-C₄ hydroxyalkyl, halo, CN, -
NR⁶R⁷, NR⁹COR¹⁰, -NR⁶S(O)_nR⁷, S(O)_nNR⁶R⁷, C₁-
C₄ haloalkyl, -OR⁷, SH or -S(O)_nR¹²;

25 R³ is selected from NR^{6a}R^{7a} and OR⁷;

R⁴ is independently selected at each occurrence from:
C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl,
30 C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, NO₂,
halo, CN, C₁-C₄ haloalkyl, NR⁶R⁷, NR⁸COR⁷,
NR⁸CO₂R⁷, COR⁷, OR⁷, CONR⁶R⁷, CO(NOR⁹)R⁷, CO₂R⁷,
or S(O)_nR⁷, where each such C₁-C₁₀ alkyl, C₂-
C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl
35 and C₄-C₁₂ cycloalkylalkyl are optionally
substituted with 1 to 3 substituents

independently selected at each occurrence from C₁-C₄ alkyl, N⁰₂, halo, CN, NR⁶R⁷, NR⁸COR⁷, NR⁸CO₂R⁷, COR⁷ OR⁷, CONR⁶R⁷, CO₂R⁷, CO(NOR⁹)R⁷, or S(O)_nR⁷;

5

R⁶, R⁷, R^{6a} and R^{7a} are independently selected at each occurrence from:

-H,
-C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl,

10 C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each optionally substituted with 1 to 3

15 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl, -aryl, aryl(C₁-C₄ alkyl), heteroaryl, heteroaryl(C₁-C₄ alkyl), heterocyclyl or heterocyclyl(C₁-C₄ alkyl);

25

alternatively, NR⁶R⁷ and NR^{6a}R^{7a} are independently piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine or thiomorpholine, each optionally substituted with 1-3 C₁-C₄ alkyl groups;

30

R⁸ is independently selected at each occurrence from H or C₁-C₄ alkyl;

35 R⁹ and R¹⁰ are independently selected at each occurrence from H, C₁-C₄ alkyl, or C₃-C₆ cycloalkyl;

R¹¹ is selected from H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, or C₃-C₆ cycloalkyl;

5 R¹² is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

R¹³ is selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, aryl, aryl(C₁-C₄ alkyl)-, 10 heteroaryl or heteroaryl(C₁-C₄ alkyl)-;

R¹⁵ and R¹⁶ are independently selected at each occurrence from H, C₁-C₆ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₆ cycloalkylalkyl, except that 15 for S(O)_nR¹⁵, R¹⁵ cannot be H;

aryl is phenyl or naphthyl, each optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

25 heteroaryl is pyridyl, pyrimidinyl, triazinyl, furanyl, pyranyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, 2,3-dihydrobenzothienyl or 2,3-dihydrobenzofuranyl, 30 each being optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, -COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵,

35

$N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{16}R^{15}$, and
 $CONR^{16}R^{15}$;

heterocyclyl is saturated or partially saturated
5 heteroaryl, optionally substituted with 1 to 5
substituents independently selected at each
occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl,
halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH,
S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵,
10 $N(COR^{15})_2$, $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{15}$, $NR^{15}R^{16}$, and
 $CONR^{16}R^{15}$;


n is independently at each occurrence 0, 1 or 2.

15 2. A method of claim 1 wherein, in the compound of
Formula (1), Ar is phenyl, pyridyl or 2,3-
dihydrobenzofuranyl, each optionally substituted with
1 to 4 R⁴ substituents.

20 3. A method of claim 1 wherein, in the compound of
Formula (1), Ar is 2,4-dichlorophenyl, 2,4-
dimethylphenyl or 2,4,6-trimethylphenyl, R¹ and R² are
CH₃, and R³ is NR^{6a}R^{7a}.

25 4. A compound of Formula (1):

254



(1)

and isomers thereof, stereoisomeric forms thereof, or
 5 mixtures of stereoisomeric forms thereof, and
 pharmaceutically acceptable salt forms thereof
 wherein:

Ar is selected from phenyl, naphthyl, pyridyl,
 10 pyrimidinyl, triazinyl, furanyl, thienyl,
 benzothienyl, benzofuranyl, 2,3-
 dihydrobenzofuranyl, 2,3-dihydrobenzothienyl,
 indanyl, 1,2-benzopyranyl, 3,4-dihydro-1,2-
 benzopyranyl, tetralinyl, each Ar optionally
 15 substituted with 1 to 5 R⁴ groups and each Ar is
 attached to an unsaturated carbon atom;

R¹ is independently selected at each occurrence from
 H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl,
 20 halo, CN, C₁-C₄ haloalkyl, C₁-C₁₂ hydroxyalkyl,
 C₂-C₁₂ alkoxyalkyl, C₂-C₁₀ cyanoalkyl, C₃-C₆
 cycloalkyl, C₄-C₁₀ cycloalkylalkyl, NR⁹R¹⁰, C₁-
 C₄ alkyl-NR⁹R¹⁰, NR⁹COR¹⁰, OR¹¹, SH or S(O)_nR¹²;

25 R² is selected from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-
 C₄ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₀
 cycloalkylalkyl, C₁-C₄ hydroxyalkyl, halo, CN, -

NR⁶R⁷, NR⁹COR¹⁰, -NR⁶S(O)_nR⁷, S(O)_nNR⁶R⁷, C₁-C₄ haloalkyl, -OR⁷, SH or -S(O)_nR¹²;

R³ is selected from NR^{6a}R^{7a} and OR⁷;

5

R⁴ is independently selected at each occurrence from:

C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl,

C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, NO₂,

halo, CN, C₁-C₄ haloalkyl, NR⁶R⁷, NR⁸COR⁷,

10 NR⁸CO₂R⁷, COR⁷, OR⁷, CONR⁶R⁷, CO(NOR⁹)R⁷, CO₂R⁷, or S(O)_nR⁷, where each such C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl and C₄-C₁₂ cycloalkylalkyl are optionally substituted with 1 to 3 substituents

15

independently selected at each occurrence from C₁-C₄ alkyl, NO₂, halo, CN, NR⁶R⁷, NR⁸COR⁷, NR⁸CO₂R⁷, COR⁷ OR⁷, CONR⁶R⁷, CO₂R⁷, CO(NOR⁹)R⁷, or S(O)_nR⁷;

20

R⁶, R⁷, R^{6a} and R^{7a} are independently selected at each occurrence from:

-H,

-C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl, C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈

25

alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each

optionally substituted with 1 to 3

30

substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl,

cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵,

NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl,

35

heteroaryl or heterocyclyl,

000250020060
000250020060

-aryl, aryl(C₁-C₄ alkyl), heteroaryl,
heteroaryl(C₁-C₄ alkyl), heterocyclyl or
heterocyclyl(C₁-C₄ alkyl),
alternatively, NR⁶R⁷ and NR^{6a}R^{7a} are independently
5 piperidine, pyrrolidine, piperazine, N-
methylpiperazine, morpholine or thiomorpholine, each
optionally substituted with 1-3 C₁-C₄ alkyl groups;

R⁸ is independently selected at each occurrence from H
10 or C₁-C₄ alkyl;

R⁹ and R¹⁰ are independently selected at each
occurrence from H, C₁-C₄ alkyl, or C₃-C₆
cycloalkyl;

15 R¹¹ is selected from H, C₁-C₄ alkyl, C₁-C₄ haloalkyl,
or C₃-C₆ cycloalkyl;

R¹² is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

20 R¹³ is selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-
C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-
C₁₂ cycloalkylalkyl, aryl, aryl(C₁-C₄ alkyl)-,
heteroaryl or heteroaryl(C₁-C₄ alkyl)-;

25 R¹⁵ and R¹⁶ are independently selected at each
occurrence from H, C₁-C₆ alkyl, C₃-C₁₀
cycloalkyl, C₄-C₁₆ cycloalkylalkyl, except that
for S(O)_nR¹⁵, R¹⁵ cannot be H;

30 aryl is phenyl or naphthyl, each optionally
substituted with 1 to 5 substituents
independently selected at
each occurrence from C₁-C₆ alkyl, C₃-
C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano,

35

OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

5 heteroaryl is pyridyl, pyrimidinyl, triazinyl, furanyl, pyranyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, 2,3-dihydrobenzothienyl or 2,3-dihydrobenzofuranyl, each being optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, -COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

20 heterocyclyl is saturated or partially saturated heteroaryl, optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁵R¹⁶, and CONR¹⁶R¹⁵;

25 n is independently at each occurrence 0, 1 or 2; with the provisos that:

30 (1) when R² is H and R³ is -OR⁷ and R⁷ is H, then R¹ is not H, OH or SH;

(2) when R^1 is CH_3 or C_2H_5 and R^2 is H, and R^3 is OH, NHC_4H_9 , or $N(C_2H_5)_2$, then Ar is not phenyl or m- CH_3 -phenyl;

5 (3) when R^2 is H and Ar is pyridyl, pyrimidinyl or pyrazinyl, and R^3 is $NR^{6a}R^{7a}$, then R^{6a} and R^{7a} are not H or alkyl;

(4) when R^2 is $SO_2NR^6R^7$, then R^3 is not OH; and

10 (5) when R^2 is $-NR^6SO_2R^7$ or $-SO_2NR^6R^7$, then R^3 is not OH.

15 2. A compound of claim ~~1~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof with the additional provisos that: (1) when R^1 is H, C_1-C_4 alkyl, halo, CN, C_1-C_{12} hydroxyalkyl, C_1-C_4 alkoxyalkyl or $SO_2(C_1-C_4$ alkyl) and R^3 is $NR^{6a}R^{7a}$ and R^{6a} is unsubstituted C_1-C_4 alkyl, then R^{7a} is not phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, furanyl, benzofuranyl, benzothiazolyl, indolyl or C_3-C_6 cycloalkyl; and (2) when R^1 is H, C_1-C_4 alkyl, halo, CN, C_1-C_{12} hydroxyalkyl, C_1-C_4 alkoxyalkyl or $SO_2(C_1-C_4$ alkyl) and R^3 is $NR^{6a}R^{7a}$ and R^{7a} is unsubstituted C_1-C_4 alkyl, then R^{6a} is not phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, furanyl, benzofuranyl, benzothiazolyl, indolyl or C_3-C_6 cycloalkyl.

3. A compound of claim ~~1~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein: Ar is phenyl,

pyridyl or 2,3-dihydrobenzofuranyl, each optionally substituted with 1 to 4 R⁴ substituents.

4 3
7. A compound of claim 8 and isomers thereof,
5 stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein: Ar is 2,4-dichlorophenyl, 2,4-dimethylphenyl or 2,4,6-trimethylphenyl, and R¹ and R² are CH₃.

10 5
8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 4.

15 6
9. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 6.

20 7
10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 4.

25 8 3
11. A compound of claim 6 and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:
R^{6a} is independently selected from:
-H,
-C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl,
C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each
optionally substituted with 1 to 3 substituents independently selected at each

occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl, -aryl, aryl(C₁-C₄ alkyl)-, heteroaryl, heteroaryl(C₁-C₄ alkyl)-, heterocyclyl or heterocyclyl(C₁-C₄ alkyl)-; and

10 R^{7a} is independently selected at each occurrence from: -H, -C₅-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl, C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each 15 optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl, -aryl, aryl(C₁-C₄ alkyl), heteroaryl, heteroaryl(C₁-C₄ alkyl), heterocyclyl or heterocyclyl(C₁-C₄ alkyl);

20 alternatively, NR⁶R⁷ and NR^{6a}R^{7a} are independently piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine or thiomorpholine, each 25 optionally substituted with 1-3 C₁-C₄ alkyl groups.

30 12. A compound of claim 9 and isomers thereof, 35 stereoisomeric forms thereof, or mixtures of

261

stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

5 R^{6a} and R^{7a} are identical and are selected from:
-C₁-C₄ alkyl or C₃-C₆ cycloalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, -COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl, and -aryl or heteroaryl.

10 15 ¹⁰ 13. A compound of claim 6 and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

20 R^{6a} is selected from:
-H,
-C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl,
C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl,

25 30

-aryl, aryl(C₁-C₄ alkyl), heteroaryl,
heteroaryl(C₁-C₄ alkyl), heterocyclyl or
heterocyclyl(C₁-C₄ alkyl);

R^{7a} is selected from:

5 -C₁-C₄ alkyl and each such C₁-C₄ alkyl is
substituted with 1-3 substituents
independently selected at each occurrence from
C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄
haloalkyl, cyano, OR¹⁵, SH, S(O)nR¹³, COR¹⁵,
10 CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂,
NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵,
aryl, heteroaryl or heterocyclyl.

11 14. A compound of claim ~~8~~³ and isomers thereof,
15 stereoisomeric forms thereof, or mixtures of
stereoisomeric forms thereof, and pharmaceutically
acceptable salt forms thereof wherein:

one of R^{6a} and R^{7a} is selected from:
20 -C₃-C₆ cycloalkyl, each such C₃-C₆ cycloalkyl
optionally substituted with 1-3 substituents
independently selected at each occurrence from
C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄
haloalkyl, cyano, OR¹⁵, SH, S(O)nR¹³, COR¹⁵,
25 CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂,
NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵,
aryl, heteroaryl or heterocyclyl,
-aryl,
-heteroaryl or
30 -heterocyclyl,
and the other of R^{6a} and R^{7a} is unsubstituted C₁-C₄
alkyl.

12 15. A compound of claim ~~8~~³ and isomers thereof,
35 stereoisomeric forms thereof, or mixtures of

868270 20007060

stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein R^{6a} and R^{7a} are independently H or C₁-C₁₀ alkyl, each such C₁-C₁₀ alkyl optionally substituted with

5 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, R⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl,

10 heteroaryl or heterocyclyl.

13

16. A compound of claim ~~14~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

15 -Ar is phenyl, pyridyl or 2,3-dihydrobenzofuranyl, and each Ar is optionally substituted with 1 to 4 R⁴ substituents;

-R¹ and R² are independently selected from H, C₁-C₄

20 alkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

14

17. A compound of claim ~~11~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of

25 stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

-Ar is phenyl, pyridyl or 2,3-dihydrobenzofuranyl, and each Ar is optionally substituted with 1 to 4 R⁴ substituents;

-R¹ and R² are independently selected from H, C₁-C₄

30 alkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

15

18. A compound of claim ~~12~~ and isomers thereof,

35 stereoisomeric forms thereof, or mixtures of

stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

-Ar is phenyl, pyridyl or 2,3-dihydrobenzofuranyl, and each Ar is optionally substituted with 1 to 4 R⁴ substituents;

5 -R¹ and R² are independently selected from H, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

10 ¹⁶ ~~19~~ ¹⁰. A compound of claim ~~13~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

15 -Ar is phenyl, pyridyl or 2,3-dihydrobenzofuranyl, and each Ar is optionally substituted with 1 to 4 R⁴ substituents;

-R¹ and R² are independently selected from H, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

20 ¹⁷ ~~20~~ ¹¹. A compound of claim ~~14~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

25 -Ar is phenyl, pyridyl or 2,3-dihydrobenzofuranyl, and each Ar is optionally substituted with 1 to 4 R⁴ substituents;

-R¹ and R² are independently selected from H, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

30 ¹⁸ ~~21~~ ¹³. A compound of claim ~~16~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein:

265

one of R^{6a} and R^{7a} is selected from:

-C₃-C₆ cycloalkyl, each such C₃-C₆ cycloalkyl
optionally substituted with 1-3 substituents
independently selected at each occurrence from
C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄
haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵,
CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂,
NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵,
aryl, heteroaryl or heterocyclyl,

-aryl,
-heteroaryl or
-heterocyclyl,

and the other of R^{6a} and R^{7a} is unsubstituted C₁-C₄ alkyl.

19 13
22. A compound of claim 16 and isomers thereof,
stereoisomeric forms thereof, or mixtures of
stereoisomeric forms thereof, and pharmaceutically
acceptable salt forms thereof wherein

20 R^{6a} and R^{7a} are independently H or C₁-C₁₀ alkyl,
each such C₁-C₁₀ alkyl optionally substituted with
1 to 3 substituents independently selected at each
occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl,
halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³,
COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂,
NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl,
heteroaryl or heterocyclyl.

20
23. A compound of claim 4 and isomers thereof,
30 stereoisomeric forms thereof, or mixtures of
stereoisomeric forms thereof, and pharmaceutically
acceptable salt forms thereof wherein R¹ is
independently selected at each occurrence from H, C₁-
C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄
haloalkyl, C₁-C₁₂ hydroxyalkyl, C₂-C₁₂ alkoxyalkyl,
C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl.

21

~~25~~ 24. A compound of claim ~~4~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein R² is selected from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl, C₁-C₄ hydroxyalkyl, halo, CN, -NR⁶R⁷, C₁-C₄ haloalkyl, -OR⁷.

22

10 ~~25~~ 25. A compound of claim ~~4~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein R⁴ is independently selected at each occurrence from: H, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, halo, CN, C₁-C₄ haloalkyl, NR⁶R⁷, COR⁷, OR⁷, S(O)_n(C₁-C₁₀ alkyl), where each such C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₃-C₆ cycloalkyl and C₄-C₁₂ cycloalkylalkyl are 20 optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₄ alkyl, NR⁶R⁷, COR⁷ OR⁷, CO₂R⁷ and where R⁷ in SONR⁷ is C₁-C₁₀ alkyl.

23


25 ~~26~~ 26. A compound of claim ~~4~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof wherein R⁴ is independently selected at each occurrence from: H, C₁-C₁₀ alkyl, C₁-C₄ alkoxy, halo, CN and -NR⁶R⁷.

26 *The compound of claim 1 which is a*

27. ~~A compound of Formula (50)~~

267

T2680



FORMULA (50)

5 and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof, selected from the group consisting of:

10

a compound of Formula (50) wherein R³ is -NHCH(n-Pr)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

15

a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

20

a compound of Formula (50) wherein R³ is -NHCH(Et)(n-Bu), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

25

a compound of Formula (50) wherein R³ is -NHCH(Et)(CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

30

a compound of Formula (50) wherein R³ is -NHCH(CH₂OEt)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

268

a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{Et})_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $-\text{N}(\text{Me})(\text{Ph})$, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{Et})(\text{n-Pr})$, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{CH}_2\text{OMe})_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is Me;

20 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{CH}_2\text{OMe})_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{Et})(\text{CH}_2\text{OMe})$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{Et})_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $-\text{OEt}$, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $-\text{N}(\text{Et})_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

45 a compound of Formula (50) wherein R^3 is $-\text{N}(\text{CH}_2\text{CN})_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

50 a compound of Formula (50) wherein R^3 is $-\text{NHCH}(\text{Me})(\text{CH}_2\text{OMe})$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

00000000000000000000000000000000

a compound of Formula (50) wherein R³ is -
 5 OCH(Et)(CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Me,
 R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(n-
 10 Pr)(CH₂cPr), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is
 H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is -
 NHCH(Me)(CH₂N(Me)₂), R^{4a} is Me, R^{4b} is H, R^{4c} is
 Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
 20 N(cPr)(CH₂CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Me,
 R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(n-
 25 Pr)(CH₂CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d}
 is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(n-
 30 Bu)(CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is
 H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
 35 NHCH(Et)(CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Me,
 R^{4d} is H and R^{4e} is Me;

a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a}
 40 is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is Me;

a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂,
 R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is
 Me;

a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)₂,
 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e}
 is H;

a compound of Formula (50) wherein R^3 is -
 $NHCH(Et)(CH_2OMe)$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe,
 R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $-N(Et)_2$, R^{4a} is
Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is Me;

a compound of Formula (50) wherein R^3 is $-NHCH(CH_2OEt)_2$,
10 R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is
Me;

15 a compound of Formula (50) wherein R^3 is -
 $NHCH(CH_2CH_2OMe)(CH_2OMe)$, R^{4a} is Me, R^{4b} is H, R^{4c}
is Me, R^{4d} is H and R^{4e} is Me;

20 a compound of Formula (50) wherein R^3 is morpholino, R^{4a}
is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $-N(CH_2CH_2OMe)_2$,
25 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e}
is H;

a compound of Formula (50) wherein R^3 is $-NHCH(Et)_2$, R^{4a}
is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $-NH(c-Pr)$, R^{4a}
is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $-NHCH(CH_2OMe)_2$,
35 R^{4a} is CN, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e}
is H;

a compound of Formula (50) wherein R^3 is $-N(c-$
 $Pr)(CH_2CH_2CN)$, R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d}
is H and R^{4e} is Me;

40 a compound of Formula (50) wherein R^3 is $-NCH(CH_2OMe)_2$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is Br, R^{4d} is H and R^{4e} is
H;

a compound of Formula (50) wherein R³ is -
 NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Br, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is -NHCH(Et)(CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is -N(C-Pr)(CH₂CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is -N(C-Pr)(CH₂CH₂CN), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is (S)-NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Br, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
NH(CH₂OMe)(CH₂-iPr), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is H, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is NMe₂, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is -
NHCH(CH₂OMe)(n-Pr), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
NHCH(CH₂OEt)(Et), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is -
NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is NMe₂, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -
NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is Br, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

10 10 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is NMe₂, R^{4d} is H and R^{4e} is H;

15 15 a compound of Formula (50) wherein R³ is (S)-NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Me, R^{4d} is H and R^{4e} is H;

20 20 a compound of Formula (50) wherein R³ is (S)-NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is (S)-NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

25 25 a compound of Formula (50) wherein R³ is (S)-NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

30 30 a compound of Formula (50) wherein R³ is -NHCH(CH₂OMe)(CH₂CH₂OMe), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(c-Pr)(CH₂CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

35 35 a compound of Formula (50) wherein R³ is -NH(Et)(CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

40 40 a compound of Formula (50) wherein R³ is -NH(Et)(CH₂CN), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Me, R^{4b} is Me, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is -
N(CH₂CH₂OMe)(CH₂CH₂OH), R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is -N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is Me, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is Me, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is -N(c-Pr) (n-Pr), R^{4a} is Me, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is -NHCH (Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is -NHCH(Et) (CH₂OMe), R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is -N(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is CN, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is -N(c-Pr) (CH₂CH₂CN), R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is -NHCH(CH₂OH)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is Cl, R^{4d} is H and R^{4e} is H; and

5 a compound of Formula (50) wherein R³ is -NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is 2-ethylpiperid-1-yl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is cyclobutyl-amino, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(Me)CH₂CH=CH₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(Pr)CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is N(Me)Bu, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is N(Me)propargyl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is N(Et)propargyl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Et, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Pr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is NHCH(cPr)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is 2-ethylpiperid-1-yl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

277

a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $N(Et)CH_2CH=CH_2$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $N(Me)CH_2cPr$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $N(Et)CH_2cPr$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $N(Pr)CH_2cPr$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $N(Me)Et$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $N(Me)Bu$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Et)propargyl$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

228

a compound of Formula (50) wherein R^3 is
 $NHCH(CH_3)CH(CH_3)CH_3$, R^{4a} is Me, R^{4b} is H, R^{4c} is
 OMe , R^{4d} is Me and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-$
 $CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is
Me and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Me$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me and R^{4e}
is H;

15 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Et$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me and R^{4e}
is H;

20 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me and R^{4e}
is H;

25 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-$
 CH_2cPr , R^{4a} is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me
and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $-NHCH(Et)_2$, R^{4a}
is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me and R^{4e} is
H;

35 a compound of Formula (50) wherein R^3 is $NHCH(cPr)_2$, R^{4a}
is Me, R^{4b} is H, R^{4c} is OMe , R^{4d} is Me and R^{4e} is
H;

40 a compound of Formula (50) wherein R^3 is $-NHCH(Et)_2$, R^{4a}
is OMe , R^{4b} is H, R^{4c} is OMe , R^{4d} is H and R^{4e} is
H;

a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $N(Et)CH_2CH=CH_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $N(Me)CH_2cPr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $N(Et)CH_2cPr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $N(Pr)CH_2cPr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $N(Me)Pr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Me)Et$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is N(Et)propargyl, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Et, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Pr, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH₂CH₃, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is NHCH(cPr)₂, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)₂, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(Et)_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $N(Et)CH_2CH=CH_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $N(Et)CH_2cPr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $N(Pr)CH_2cPr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $N(Me)Pr$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(Me)Et$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Me)Bu$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(Me)propargyl$,
 R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

5 a compound of Formula (50) wherein R^3 is $N(Et)propargyl$,
 R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

10 a compound of Formula (50) wherein R^3 is
 $NHCH(CH_3)CH(CH_3)CH_3$, R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

15 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-$
 $CH_2CH=CH_2$, R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

20 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Me$,
 R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

25 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$,
 R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

30 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-$
 CH_2cPr , R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

35 a compound of Formula (50) wherein R^3 is
 $NHCH(CH_3)CH_2CH_3$, R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe ,
 R^{4d} is Me and R^{4e} is H ;

40 a compound of Formula (50) wherein R^3 is $NHCH(cPr)_2$, R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)_2$,
 R^{4a} is OMe , R^{4b} is H , R^{4c} is OMe , R^{4d} is Me and R^{4e} is H ;

283 ~~284~~ ~~285~~

a compound of Formula (50) wherein R^3 is $NHCH(Et)_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(Et)_2$, R^{4a} is OMe, R^{4b} is H, R^{4c} is OMe, R^{4d} is Me and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

15 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

20 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

25 a compound of Formula (50) wherein R^3 is $N(Et)CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

30 a compound of Formula (50) wherein R^3 is $N(Me)CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

35 a compound of Formula (50) wherein R^3 is $N(Pr)CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

40 a compound of Formula (50) wherein R^3 is $N(Me)Pr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

a compound of Formula (50) wherein R^3 is $N(Me)Et$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

a compound of Formula (50) wherein R³ is N(Me)Bu, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

5 a compound of Formula (50) wherein R³ is N(Me)propargyl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

10 a compound of Formula (50) wherein R³ is N(Et)propargyl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

15 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

20 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

25 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Et, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

30 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Pr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

35 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

40 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH₂CH₃, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

5 a compound of Formula (50) wherein R³ is NHCH(cPr)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is Me;

10 a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

15 a compound of Formula (50) wherein R³ is 2-ethylpiperid-1-yl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

20 a compound of Formula (50) wherein R³ is N(Me)CH₂CH=CH₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

30 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

35 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

40 a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

~~285~~ 286

a compound of Formula (50) wherein R^3 is $N(Me)Bu$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

5 a compound of Formula (50) wherein R^3 is $N(Me)propargyl$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

10 a compound of Formula (50) wherein R^3 is $N(Et)propargyl$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

15 a compound of Formula (50) wherein R^3 is $NHCH(CH_3)CH(CH_3)CH_3$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

20 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Et$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

30 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

35 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

40 a compound of Formula (50) wherein R^3 is $NHCH(cPr)_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

5 a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

10 a compound of Formula (50) wherein R³ is N(Et)₂, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

15 a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

20 a compound of Formula (50) wherein R³ is 2-ethylpiperid-1-yl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R³ is N(Me)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

30 a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

35 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

40 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

5 a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

a compound of Formula (50) wherein R³ is N(Me)Bu, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

10 a compound of Formula (50) wherein R³ is N(Me)propargyl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

15 a compound of Formula (50) wherein R³ is N(Et)propargyl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

20 a compound of Formula (50) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

30 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Et, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

35 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Pr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

40 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

a compound of Formula (50) wherein R³ is NHCH(CH₃)CH₂CH₃, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

5 a compound of Formula (50) wherein R³ is NHCH(cPr)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

10 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

15 a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

20 a compound of Formula (50) wherein R³ is N(Et)₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is OMe;

25 a compound of Formula (50) wherein R³ is 2-ethylpiperid-1-yl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is cyclobutyl-amino, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R³ is N(Me)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

201 290

060210-2005T060

a compound of Formula (50) wherein R³ is N(Et)CH₂cPr,
R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is N(Pr)CH₂cPr,
R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(Me)propargyl,
R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(Et)propargyl,
R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is
NHCH(CH₃)CH(CH₃)CH₃, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-
CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me,
R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2cPr$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $NHCH(CH_3)CH_2CH_3$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $NHCH(cPr)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $N(Et)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is H and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(Pr)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(Me)Bu, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is NH(CH(CH₃)CH(CH₃)CH₃), R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

2Mf 293

a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Et$,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2cPr$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $NH(CH(CH_3)CH_2CH_3)$, R^{4a} is Cl, R^{4b} is F, R^{4c} is OMe,
 R^{4d} is H and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $NHCH(cPr)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

~~25~~ a compound of Formula (50) wherein R^3 is $N(Et)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H.

30 a compound of Formula (50) wherein R^3 is $NHCH(Et)_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is N(Et)CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(Pr)CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(Me)Pr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(Me)propargyl, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is NH(CH(CH₃)CH(CH₃)CH₃), R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

45 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

~~2016~~ 295

a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Me,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Et,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)Pr,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)-
 CH₂cPr, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is
 NHCH(CH₃)CH₂CH₃, R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe,
 R^{4d} is OMe and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is NHCH(cPr)₂, R^{4a}
 is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(CH₂CH₂OMe)₂,
 R^{4a} is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(Et)₂, R^{4a}
 is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H.

a compound of Formula (50) wherein R³ is NHCH(Et)₂, R^{4a}
 is Cl, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is 2-ethylpiperid-
 1-yl, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is cyclobutyl-amino, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is N(Me)CH₂CH=CH₂, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

15 a compound of Formula (50) wherein R³ is N(Me)CH₂cPr, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

20 a compound of Formula (50) wherein R³ is N(Et)CH₂cPr, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

25 a compound of Formula (50) wherein R³ is N(Pr)CH₂cPr, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

30 a compound of Formula (50) wherein R³ is N(Me)Et, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R³ is N(Me)Bu, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R³ is NH(CH(CH₃)CH(CH₃)CH₃), R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2CH=CH_2$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

5 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Me$,
 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

10 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Et$,
 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

15 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)Pr$,
 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

20 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)-CH_2cPr$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

25 a compound of Formula (50) wherein R^3 is $NH(CH(CH_3)CH_2CH_3)$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

30 a compound of Formula (50) wherein R^3 is $N(CH_2CH_2OMe)_2$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $NHCH(Et)_2$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 ~~a compound of Formula (50) wherein R^3 is $N(Et)_2$, R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H.~~

~~2009~~ 2088

10 a compound of Formula (50) wherein R^3 is 2-ethylpiperid-1-yl, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

5 15 a compound of Formula (50) wherein R^3 is cyclobutyl-amino, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

10 20 a compound of Formula (50) wherein R^3 is $N(Me)CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

15 25 a compound of Formula (50) wherein R^3 is $N(Et)CH_2CH=CH_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

20 30 a compound of Formula (50) wherein R^3 is $N(Me)CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is F and R^{4e} is H;

25 35 a compound of Formula (50) wherein R^3 is $N(Et)CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

30 40 a compound of Formula (50) wherein R^3 is $N(Pr)CH_2cPr$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

35 a compound of Formula (50) wherein R^3 is $N(Me)Et$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Me)Bu$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

40 a compound of Formula (50) wherein R^3 is $N(Me)propargyl$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H;

280 299

a compound of Formula (50) wherein R³ is
 $\text{NH}(\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_3$, R^{4a} is Br, R^{4b} is H, R^{4c} is
 OMe, R^{4d} is OMe and R^{4e} is H;

5 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})-$
 $\text{CH}_2\text{CH}=\text{CH}_2$, R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is
 F and R^{4e} is H;

10 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})\text{Me}$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e}
 is H;

15 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})\text{Et}$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e}
 is H;

20 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})\text{Pr}$,
 R^{4a} is Br, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e}
 is H;

25 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})-$
 CH_2cPr , R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe
 and R^{4e} is H;


30 a compound of Formula (50) wherein R³ is $\text{NHCH}(\text{cPr})_2$, R^{4a}
 is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is
 H;

35 a compound of Formula (50) wherein R³ is $\text{N}(\text{CH}_2\text{CH}_2\text{OMe})_2$,
 R^{4a} is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e}
 is H;

40 a compound of Formula (50) wherein R³ is $\text{NHCH}(\text{Et})_2$, R^{4a}
 is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is
 H; and

a compound of Formula (50) wherein R³ is $\text{N}(\text{Et})_2$, R^{4a}
 is Me, R^{4b} is H, R^{4c} is OMe, R^{4d} is OMe and R^{4e} is H.

25
28. A compound of claim 4 of Formula (60)



FORMULA (60)

5

and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof, selected from the group consisting of:

10

a compound of Formula (60) wherein R^3 is $NHCH(Et)_2$, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

15

a compound of Formula (60) wherein R^3 is 2-ethylpiperid-1-yl, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is cyclobutyl-amino, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

20

a compound of Formula (60) wherein R^3 is $N(Me)CH_2CH=CH_2$, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(Et)CH_2cPr$, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

25

a compound of Formula (60) wherein R^3 is $N(Pr)CH_2cPr$, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

100-20057060

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is NH(CH(CH₃)CH₂CH₃), Ar is 6-dimethylamino-4-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is NHCH(cPr)₂ Ar is 6-dimethylamino-4-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is NHCH(Et)₂ Ar is 6-dimethylamino-4-methylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(Et)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is 2-ethylpiperid-1-yl, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is cyclobutyl-amino, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)CH₂CH=CH₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Et)CH₂cPr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Pr)CH₂cPr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is N(Me)Pr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)Et, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is N(Me)Bu, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)propargyl, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is NH(CH(CH₃)CH(CH₃)CH₃), Ar is 6-dimethylamino-4-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Me,

25 Ar is 6-dimethylamino-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Et, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Pr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

169270-20057060

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is NH(CH(CH₃)CH₂CH₃), Ar is 6-dimethylamino-4-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is NHCH(cPr)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is NHCH(Et)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(Et)₂, Ar is 6-dimethylamino-4-methylpyrid-3-yl.

30 a compound of Formula (60) wherein R³ is 2-ethylpiperid-1-yl, Ar is 6-methoxy-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is cyclobutyl-amino, Ar is 6-methoxy-4-methylpyrid-3-yl;

35 a compound of Formula (60) wherein R³ is N(Me)CH₂CH=CH₂, Ar is 6-methoxy-4-methylpyrid-3-yl;

40 a compound of Formula (60) wherein R³ is N(Et)CH₂cPr, Ar is 6-methoxy-4-methylpyrid-3-yl;

45 a compound of Formula (60) wherein R³ is N(Pr)CH₂cPr,

Ar is 6- methoxy -4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)Pr, Ar is
6- methoxy -4-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is N(Me)Et, Ar is
6- methoxy -4-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is N(Me)Bu, Ar is
6- methoxy -4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)propargyl,
Ar is 6- methoxy -4-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is N(Et)propargyl,
Ar is 6- methoxy -4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is
NHCH(CH₃)CH(CH₃)CH₃, Ar is 6- methoxy -4-
20 methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-
CH₂CH=CH₂, Ar is 6- methoxy -4-methylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Me,
Ar is 6- methoxy -4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Et,
Ar is 6- methoxy -4-methylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Pr,
Ar is 6- methoxy -4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, Ar is 6-methoxy-4-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is NHCH(CH₃)CH₂CH₃, Ar is 6-methoxy-4-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is NHCH(cPr)₂ Ar is 6-methoxy-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)₂, Ar is 6-methoxy-4-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is NHCH(Et)₂ Ar is 6-methoxy-4-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is N(Et)₂, Ar is 6-methoxy-4-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is 2-ethylpiperid-1-yl, Ar is 4-methoxy-6-methylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is cyclobutyl-amino, Ar is 4-methoxy-6-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)CH₂CH=CH₂, Ar is 4-methoxy-6-methylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(Et)CH₂cPr, Ar is 4-methoxy-6-methylpyrid-3-yl;

368270-20057060

a compound of Formula (60) wherein R³ is N(Pr)CH₂cPr, Ar is 4-methoxy-6-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is N(Me)Pr, Ar is 4-methoxy-6-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)Et, Ar is 4-methoxy-6-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is N(Me)Bu, Ar is 4-methoxy-6-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Me)propargyl, Ar is 4-methoxy-6-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, Ar is 4-methoxy-6-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, Ar is 4-methoxy-6-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Me, Ar is 4-methoxy-6-methylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Et, Ar is 4-methoxy-6-methylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Pr, Ar is 4-methoxy-6-methylpyrid-3-yl;

30

2682005T060

a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, Ar is 4-methoxy-6-methylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is NH(CH(CH₃)CH₂CH₃), Ar is 4-methoxy-6-methylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is NHCH(cPr)₂, Ar is 4-methoxy-6-methylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is NHCH(Et)₂, Ar is 6-methoxy-4-methylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is 2-ethylpiperid-1-yl, Ar is 4,6-dimethylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is cyclobutyl-amino, Ar is 4,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(Me)CH₂CH=CH₂, Ar is 4,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Et)CH₂cPr, Ar is 4,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R³ is N(Pr)CH₂cPr,

Ar is 4,6-dimethylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is N(Me)Pr, Ar is 4,6-dimethylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is N(Me)Et Ar is 4,6-dimethylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is N(Me)Bu, Ar is 4,6-dimethylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is N(Me)propargyl, Ar is 4,6-dimethylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(Et)propargyl, Ar is 4,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is NHCH(CH₃)CH(CH₃)CH₃, Ar is 4,6-dimethylpyrid-3-yl;

35 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂CH=CH₂, Ar is 4,6-dimethylpyrid-3-yl;

40 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Me, Ar is 4,6-dimethylpyrid-3-yl;

45 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Et, Ar is 4,6-dimethylpyrid-3-yl;

50 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)Pr, Ar is 4,6-dimethylpyrid-3-yl;

100057060
a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr, Ar is 4,6-dimethylpyrid-3-yl;

5 a compound of Formula (60) wherein R³ is
NHCH(CH₃)CH₂CH₃, Ar is 4,6-dimethylpyrid-3-yl;

10 a compound of Formula (60) wherein R³ is NHCH(cPr)₂, Ar
is 4,6-dimethylpyrid-3-yl;

15 a compound of Formula (60) wherein R³ is NHCH(Et)₂ Ar is
4,6-dimethylpyrid-3-yl;

20 a compound of Formula (60) wherein R³ is N(Et)₂, Ar is
4,6-dimethylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is 2-ethylpiperid-1-yl, Ar is 2,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is cyclobutyl-amino, Ar is 2,6-dimethylpyrid-3-yl;

25 a compound of Formula (60) wherein R³ is N(Me)CH₂CH=CH₂,
Ar is 2,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(Et)CH₂cPr, Ar
is Ar is 2,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(Pr)CH₂cPr, Ar
is Ar is 2,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(Me)Pr$, Ar is 2,6-dimethylpyrid-3-yl;

5 a compound of Formula (60) wherein R^3 is $N(Me)Et$, Ar is
2,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(Me)Bu$, Ar is 2,6-dimethylpyrid-3-yl;

10

a compound of Formula (60) wherein R^3 is $N(Me)propargyl$,
 Ar is 2,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is

15 NH (CH (CH₃) CH (CH₃) CH₃), Ar is 2, 6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(CH_2CH_2OMe)-CH_2CH=CH_2$, Ar is 2,6-dimethylpyrid-3-yl;

20 a compound of Formula (60) wherein R^3 is $N(CH_2CH_2OMe)Me$,
 Ar is 2,6-dimethylpyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(CH_2CH_2OMe)Et$,
 Ar is 2,6-dimethylpyrid-3-yl;

25

a compound of Formula (60) wherein R^3 is $N(CH_2CH_2OMe)Pr$,
 Ar is 2,6-dimethylpyrid-3-yl;

30 a compound of Formula (60) wherein R³ is N(CH₂CH₂OMe)-CH₂cPr Ar is 2,6-dimethylpyrid-3-yl;

~~3/12~~ 31

a compound of Formula (60) wherein R^3 is

NH(CH(CH₃)CH₂CH₃), Ar is 2,6-dimethyl pyrid-3-yl;

a compound of Formula (60) wherein R^3 is $NHCH(cPr)_2$, Ar

5 is 2,6-dimethyl pyrid-3-yl;

a compound of Formula (60) wherein R^3 is $N(CH_2CH_2OMe)_2$,

Ar is 2,6-dimethylpyrid-3-yl;

10 a compound of Formula (60) wherein R^3 is $NHCH(Et)_2$, Ar is 2,6-dimethyl-pyrid-3-yl; and

a compound of Formula (60) wherein R^3 is $N(Et)_2$, Ar is 2,6-dimethyl-pyrid-3-yl.

15

26
29. A compound of claim ~~4~~ and isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt forms thereof, wherein said compound is selected from the group consisting of:

4-((2-butyl)amino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

25 4-((2-butyl)amino)-2,7-dimethyl-8-(2,5-di methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

4-((3-pentyl)amino)-2,7-dimethyl-8-(2,5-dimethyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

30

4-((3-pentyl)amino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

~~243~~ 312

4- (N-cyclopropylmethyl-N-propylamino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

5 4- (N-cyclopropylmethyl-N-propylamino)-2,7-dimethyl-8-(2,5-dimethyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

10 4- (N-allyl-N- (2-methoxyethyl) amino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

15 4- (N-allyl-N- (2-methoxyethyl) amino)-2,7-dimethyl-8-(2,5-dimethyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

20 4- (diallylamino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine;

25 4- (N-ethyl-N- (2-methoxyethyl) amino)-2,7-dimethyl-8-(2-methyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine; and

30 4- (N-ethyl-N- (2-methoxyethyl) amino)-2,7-dimethyl-8-(2,5-dimethyl-4-methoxyphenyl)-[1,5-a]-pyrazolo-1,3,5-triazine.

35 ³⁷
³⁸ 30. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claims ³ ¹³ ¹⁴ ²⁵ ²⁶ ²⁷, ²⁸ and ²⁹.

31. A method of treating affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in mammals comprising administering to the mammal a therapeutically effective amount of a compound of claim claims 4, 6, 11, 16, 27, 28 and 29.

20

A handwritten signature consisting of a stylized 'H' and 'A' above a stylized 'B'.A handwritten signature consisting of a stylized 'a' and 's' above a stylized 'B'.